



106

216

John A. Hargrave

Trinit


1834

*The Bancroft Library*

University of California • Berkeley

FROM THE PAPERS OF

Prof. Edmund Pinney



Digitized by the Internet Archive  
in 2008 with funding from  
Microsoft Corporation





A SHORT  
PRACTICAL TREATISE  
ON  
SPHERICAL TRIGONOMETRY;

CONTAINING A FEW SIMPLE RULES,

BY WHICH

THE GREAT DIFFICULTIES TO BE ENCOUNTERED BY  
THE STUDENT IN THIS BRANCH OF  
MATHEMATICS ARE EFFECTUALLY OBIATED.

BY OLIVER BYRNE,

PROFESSOR OF MATHEMATICS;

AUTHOR OF "THE ELEMENTS OF EUCLID, BY COLOURS," "A TREATISE ON  
ALGEBRA," "LOGARITHMS," &c. &c.

LONDON:  
PRINTED AND PUBLISHED BY A. J. VALPY, M.A.  
RED LION COURT, FLEET STREET.

1835.





TO,

V. T. HURTADO, ESQ.,

THIS WORK

IS

MOST RESPECTFULLY DEDICATED

BY

HIS HUMBLE SERVANT,

THE AUTHOR.



## P R E F A C E.

---

IT will perhaps be wondered at, after so much has been written on this subject, that I should have the temerity to offer my mite to the notice of the public, or to imagine myself capable of throwing any light upon a branch of Mathematics that has been so much illuminated by the works of the learned. I trust, however, that when the reader has heard my explanations, he will not consider that I have engaged in a useless or unprofitable undertaking. Much indeed has been written by learned and experienced Mathematicians, but their works seem rather addressed to the proficient, than to the uninitiated student in the science. Their long formulæ, complicated rules, and demonstrations, &c., perplex rather than instruct the beginner, who perhaps, terrified at the mass of difficulties before him, gives up in disgust the study of a subject, which, treated in a simple manner, he had easily acquired. The great fault of all elementary treatises on this branch appears to be the crowding too much upon the mind

of the students, and distracting their attention with useless rules and demonstrations, which retard rather than assist their progress. This I have found by experience, both in my early studies, and in the extensive practice I have since obtained in communicating knowledge to others; and I have frequently in one hour's conversation enabled a pupil to master a subject which he had in vain attempted to acquire by the perusal of the ordinary rules. In short, I felt that a plain practical Treatise on Spherical Trigonometry, in which the pupil's attention should not be distracted from the subject before him, and in which the rules should be as simple and concise as possible, would much facilitate the acquirement of the subject. I have therefore written in the same manner in which I should have explained it by oral communication with my pupils. The formulæ which I have constructed for solving each Case in Spherical Trigonometry will, I hope, obtain the sanction of the Mathematician for their correctness, and the approbation of the student for the ease and rapidity with which they enable him to master the science.

## INTRODUCTION.

---

By the word Sphere is generally understood any circular body; but the term was appropriated by the ancients to an assemblage of circles and constellations representing their "Primum Mobile." The invention of this Sphere is ascribed to various persons, but it is evidently too remote to be traced by any authentic history. The Chinese had a knowledge of the Sphere at a very early period,\* from whom it was probably transmitted to the Chaldeans, thence into Egypt and Greece; but it

\* "Xuni, 2400 ans avant J. C. fit faire une sphère d'or enrichie de pierreries où l'on avait les sept planètes et la terre au milieu."—*Historie de La Chine, par Martin*, page 76.

was most successfully studied in the famous school of Alexandria. Here Euclid, the great geometrician, wrote a Treatise on the Sphere, entitled *The Phenomena*, which explained the most interesting parts of ancient astronomy; such as the right oblique ascension of the heavenly bodies, with the various other phenomena, which arise from the apparent diurnal revolution of the Primum Mobile. This work is supposed to be the first on the subject perfectly geometrical: it served long after as a model for other performances on the subject, and is still in existence, but very scarce.

Hipparchus, who flourished about two centuries after Euclid, and one before the Christian era, is said to have laid the foundation of Spherical Trigonometry. In succeeding ages it was improved by Ptolemy, Theodosius, and others; and much is ascribed to Geber, a learned Spaniard, who lived in the sixteenth century.

Baron Napier, however, made the most considerable improvements by his proposition of

circular parts, and his invention of Logarithms,\* and up to the present time many works of great merit have appeared in Europe. Yet the present, although the last, the author hopes will be found not the least useful.

\* The author has discovered so simple a method of constructing Logarithms, that they may be calculated with as much facility as the most trifling question in common arithmetic, thus superseding the necessity of tables.





# SPHERICAL TRIGONOMETRY.

---

## DEFINITIONS.

### I.

A sphere is a solid, such that if it be cut by a plane in any position the section will be a circle.

or,

It is formed by the revolution of a semicircle about the diameter which remains fixed.

or,

It is a solid such that all lines drawn from a point within called the centre to the surface are equal to one another.

### II.

The circles cutting the sphere are divided

into two kinds, the greater and lesser circles of the sphere: the greater passes through the centre, the less does not.

### III.

The nearest distance between any two points on the sphere, is the arc of a great circle;—for such an arc being described with the greatest radius, is less curved than the arc of any small circle.

### IV.

A spherical triangle is formed by three great circles on the surface of the sphere.

### V.

A spherical angle is formed by the inclination of two great circles on the surface of the sphere meeting in a point, called the angular point.

### VI.

Spherical triangles are also distinguished as right-angled, quadrantal and oblique:—thus

when one of the angles is  $90^\circ$  it is called right-angled.

## VII.

Any obtuse-angled spherical triangle may be divided into two right-angled spherical triangles, by letting fall from any of the angular points on the opposite side a great circle, whose plane will be perpendicular to the plane of the base.

### OF RIGHT-ANGLED SPHERICAL TRIGONOMETRY.

Right-angled triangle spherical trigonometry may be divided into the six following cases:—


- |      |                                |   |
|------|--------------------------------|---|
| I.   | When the hypotenuse and a leg, | } are given together with<br>the right angle. |
| II.  | the hypotenuse and an angle,   |   |
| III. | a leg and its opposite angle,  |   |
| IV.  | a leg and its adjacent angle,  |   |
| V.   | two legs,                      |   |
| VI.  | two angles,                    |   |

To find the remaining parts of the right-angled spherical triangle,

Place on the angular points of the annexed

five-sided figure (its construction is simple) first the base, next the perpendicular, thirdly the complement of the angle opposite the base, fourthly the complement of the hypotenuse, fifthly the complement of the angle opposite the perpendicular, and in the centre radius as in the annexed diagram.

R, Radius: B, Base: P, Perpendicular: A, Complement of the angle opposite the base: H, complement of the hypotenuse: a, Complement of the angle opposite the perpendicular.





This sign signifies that the quantities standing on the three prongs are equal.

Then we have the following general theorem.


$$\begin{array}{c}
 \begin{array}{l}
 \text{Co. Sin. B} \times \text{Co. Sin. a.} \\
 \text{Tan. P.} \times \text{Tan. H.} \\
 \text{Tan. A.} \times \text{Tan. B.} \\
 \text{Co. Sin. a.} \times \text{Co. Sin. H.} \\
 \text{Tan. a.} \times \text{Tan. P.} \\
 \text{Co. Sin. H.} \times \text{Co. Sin. a.} \\
 \text{Co. Sin. A.} \times \text{Co. Sin. P.} \\
 \text{Tan. B.} \times \text{Tan. H.}
 \end{array}
 \end{array}$$

The above formula solves all the cases in right-angled spherical trigonometry, but it is much better to solve each case separate.

## RULE FOR THE ARRANGEMENT OF ALL THE CASES.

Place in their respective places the given parts, and then refer to the general formula for the solution of the case.


[Obs.] This diagram, so constant in use, does not require Geometrical exactness, nor yet right lines, as it is only to regulate the arrangement of B. P. A. H. *a.* and R; it is no matter which of the angles we place B. in, so it be followed right or left by P. A. H. and *a.*



## CASE I.

---

WHEN THE HYPOTHENUSE AND A LEG  
ARE GIVEN TO FIND THE REST.



First suppose the leg given to be the perpendicular, find the part which is represented by  $x$  on the diagram, which stands for the complement of the angle opposite the base.

According to the formula we have  $R \times \text{Sin. } x = \text{Tan. H.} \times \text{Tan. P.}$

$$\text{Then Sin. } x = \frac{\text{Tan. H.} \times \text{Tan. P.}}{R}$$

Log. Sin.  $x = \text{Log. Tan. H.} + \text{Log. Tan. P.} - \text{Log. R.}$

Given the hypotenuse  $63^\circ 57' 7'' \therefore \text{H. } 26^\circ 2' 53''$ .


Given the perpendicular  $40^\circ 0' 0'' \text{ P.}$

The solution is as follows:

$9.6894258 + 9.9238135 - 10.0000000 = 9.6132393 = \text{Sin. } x = 24^\circ 13' 55''$  but  $x$  is the complement of the angle opposite the base, then the angle opposite the base is  $65^\circ 46' 5''$ .

Next find the part which is represented by  $x'$ , which stands for the complement of the angle opposite the perpendicular.

The solution of this first Case will be the same, if the side given was the base.



$\text{R.} \times \text{Sin. P.} = \text{Co. Sin. H.} \times \text{Co. Sin. } x'$

Consequently we have  $\text{Co. Sin. } x' = \frac{\text{R.} \times \text{Sin. P.}}{\text{Co. Sin. H.}}$

or which is the same,

$\text{Log. Co. Sin. } x' = \text{Log. R.} + \text{Sin. P.} - \text{Log. Co. Sin. H.}$



## EXAMPLE.

The solution is obtained by substituting in this formula the logarithms of R. Sin. P. and Co. Sin. H.; then:  $\text{Log. Co. Sin. } x' = 10. + 9.8080675 - 9.9534206 = 9.8546469 = 45^\circ 41' 21''$ ; but the Co. Sin. of  $x'$  is the sine of the angle opposite the perpendicular  $\therefore 45^\circ 41' 21''$  the angle required.

To find the part represented by  $x''$  which stands for the base.

$$R. \times \text{Sin. H.} = \text{Co. Sin. P.} \times \text{Co.}$$


$$\text{Sin. } x''$$

$$\text{Co. Sin. } x'' = \frac{R. \times \text{Sin. H.}}{\text{Co. Sin. P.}}$$

Then we have,  $\text{Log. Co. Sin.}$

$$x'' = \text{Log. R.} + \text{Log. Sin. H.}$$

$$- \text{Log. Co. Sin. P.}$$




## NUMERAL SOLUTION.

$\text{Log. Co. Sin. } x'' = 10. + 9.642846 - 9.884254 = 9.7585924 = 55^\circ$ . The leg represented by  $x''$ .

## CASE II.

---

WHEN THE HYPOTHENUSE AND AN AN-  
GLE ARE GIVEN TO FIND THE REMAIN-  
ING PARTS OF THE RIGHT-ANGLED  
SPHERICAL TRIANGLE.



We may either take the angle opposite the base or opposite the perpendicular, and the solution will be the same.

Let it first be required to find the value of  $x$

which stands for the complement of the angle, opposite the perpendicular.

Then according to the general formula we have  $R. \times \text{Sin. } H. = \text{Tan. } A. \times \text{Tan. } x.$

Then  $R. \times \text{Sin. H.} = \text{Tan. A.} \times \text{Tan. } x.$

$$\text{Tan. } x. = \frac{R. \times \text{Sin. H.}}{\text{Tan. A.}}$$

Then is  $\text{Log. of Tan. } x. = \text{Log. R.} + \text{Log. Sin. H.} - \text{Tan. A.}$  Let the hypotenuse be  $= 63^\circ 56' 7''$  and the given angle  $= 45^\circ 41' 21''$ .  $H. = 26^\circ 3' 53''$  or  $90^\circ - 63^\circ 56' 7''$ ;  $A. = 44^\circ 18' 39'' = 90^\circ - 45^\circ 41' 21''$ .

$$\begin{aligned} \text{Log. Tan. } x. &= 10. + 9.6428464 - 9.9895514. \\ &= 9.6532950 \end{aligned}$$

$x. = 34^\circ 13' 55''$ , the complement of angle opposite the perpendicular; the angle  $= 65^\circ 46' 5''$ .

Next let the required part be  $x'$ , which is situated in place of the base.

In each of the cases a reference to the general formula is given; and another thing may be remembered here, that the quantity required, and the two that are given, must be so arranged that two of the known quantities must be multiplied and equalled with the other known quantity;  $R.$  the radius is always known.

$$R. \times \text{Sin. } x' = \text{Co. Sin. H.} \times \text{Co. Sin. A.}$$

$$\text{Sin. } x' = \frac{\text{Co. Sin. H.} \times \text{Co. Sin. A.}}{\text{R.}} \quad \text{Therefore}$$

$$\text{Log. Sin. } x' = \text{Log. Co. Sin. H.} + \text{Log. Co. Sin. A.} - \text{Log. R.}$$

## EXAMPLES.


1. Given the hypotenuse of a right-angled spherical triangle  $45^\circ 17'$ , and the angle opposite the base  $16^\circ 27'$ , to find the base?

2. Required the base of a right-angled spherical triangle, when the hypotenuse is  $= 103^\circ 30'$ , and the angle opposite the base  $= 75^\circ 35'$ ?

Next, let it be required to find the value of  $x''$ , which stands in the diagram in the place of the perpendicular.

$\text{R.} \times \text{Sin. A.} = \text{Tan. H.} \times \text{Tan. } x''$ . Consequently, we have the value of  $x''$ , or its tangent  $= \frac{\text{R.} \times \text{Sin. A.}}{\text{Tan. H.}}$

by Logarithms, thus  
 $\text{Log. of } x'' = \text{Log. R.} + \text{Sin. of A} - \text{Log. Tan. H.}$



1. Given the hypotenuse  $63^{\circ} 53' 7''$ , and the angle opposite the perpendicular  $45^{\circ} 41' 21''$ , to find the perpendicular?

Answer  $40^{\circ}$ .

Required the perpendicular of a right-angled spherical triangle, when the hypotenuse is  $57^{\circ}$  and the angle  $30^{\circ}$ ?


## CASE III.

---

WHEN A LEG AND ITS OPPOSITE ANGLE  
ARE GIVEN TO FIND THE REMAINING  
PARTS.

Let the perpendicular and the angle opposite

it be given ; which is the same as the base, and the angle opposite the base ; for the perpendicular may be considered the base when we please. 1st.




to find the value of  $x$ , which represents the complement of the angle opposite the base.

$$R. \times \text{Sin. } a. = \text{Co. Sin. } x \times \text{Co. Sin. } P.$$

$$\text{Co. Sin. } x = \frac{R. \times \text{Sin. } a.}{\text{Co. Sin. } P.} \text{ or } \text{Log. Co. Sin. } x =$$

Log. R. + Log. Sin.  $a$ . — Log. Co. Sin. P. Suppose the perpendicular =  $55^{\circ} 0'$ , and its opposite angle =  $65^{\circ} 46' 5''$ , then we have P. =  $55^{\circ} 0' 0''$  and  $a$ . = Comp. of  $65^{\circ} 46' 5'' = 24^{\circ} 13' 55''$ . Consequently Log. Co. Sin.  $x$ . =  $10. + 9.6132407 - 9.7585913 = 9.8546494$ , the Log. Co. Sin. of  $x$ ;  $\therefore x = 45^{\circ} 41' 21''$ .

Suppose the perpendicular BC to be  $59^{\circ} 30'$ , and its opposite angle BAC  $67^{\circ} 47'$ , to find the angle opposite the base?




To find  $x'$  which stands for the other leg in the general scheme. (Page 4.)

From the general formula we have  $R. \times \text{Sin. } x' = \text{Tan. } a. \times \text{Tan. P. Sin.}$

$$x' = \frac{\text{Tan. } a. + \text{Tan. P.}}{R.} \text{ as}$$

before P. =  $55^{\circ} 0' 0''$  and  $a$ . =  $24^{\circ} 13' 55''$ . Log. Sin.  $x' = \text{Log. Tan. } a. + \text{Log. Tan. P.} - \text{Log. R.}$

$$\text{Log. Sin. } x' = 10.1547732 + 9.6532976 - 10.$$



∴ The Log. Co. Sin.  $x' = 9.8080708 = 40^\circ 0' 0''$ .

Required to find the base AC of the above right-angled spherical triangle, when the perpendicular BC is  $23^\circ 57'$ , and its opposite angle  $38^\circ 42' 3''$ .

In order to find the value of  $x$  which stands in the scheme for the complement of the hypotenuse.

$$R. \times \text{Sin. } P. = \text{Co. Sin. } a. \times \text{Co. Sin. } x''. \text{ Co.}$$

$$\text{Sin. } x'' = \frac{R. \times \text{Sin. } P.}{\text{Co. Sin. } A.}$$


$$\text{or Log. Co. Sin. } x'' = \\ \text{Log. R.} + \text{Log. Sin. } P. \\ - \text{Log. Co. Sin. } a.$$

Then as before  $P. =$

$$55^\circ 0' 0'' \quad a \quad 24^\circ 13' 55''$$

$$\text{Log. Co. Sin. } x'' = 10. + 9.9133645 - 9.9599432.$$

Log. Co. Sin.  $x'' = 9.9534213$ . Therefore  $x'' = 26^\circ 3' 53''$ . And the hypotenuse  $= 63^\circ 56' 7''$ .



Given the perpendicular BC of a right-angled spherical triangle  $= 55^\circ$ , and the angle opposite the perpendicular  $CAB = 28^\circ 15'$ , to find the hypotenuse?

Answer.



## CASE IV.

---

A LEG AND ITS ADJACENT ANGLE BEING  
GIVEN TO FIND THE REST.

Suppose the given leg =  $55^{\circ} 0' 0''$ , and its adjacent angle =  $45^{\circ} 41' 21''$ , which in the diagram stands for the perpendicular, and its adjacent angle.


$P. = 55^{\circ} 0' 0''$ ;  $A. = 44^{\circ} 18' 39''$ , the complement of the angle opposite the base.

1st, find  $x$  which stands for the base.  $R. \times \text{Sin. } P. = \text{Tan. } A. \times \text{Tan. } x$

$$\text{Tan. } x = \frac{R. \times \text{Sin. } P.}{\text{Tan. } A.}$$


$$\text{Log. Tan. } x = \text{Log. } R. + \text{Log. Sin. } P. - \text{Log.}$$

$$\text{Tan. } A. \quad \text{Log. Tan. } x = 10 + 9.9133645 - 9.$$



9895514 = 9.9238131 = Log. Tan. of  $x$ ;  $\therefore x = 40^\circ$ .

2. Suppose the given leg  $BC = 57^\circ 30'$ , and its adjacent angle  $BCA = 45^\circ 38'$ , to find the base?




Next find the part represented by  $x'$ , which stands for the complement of the angle

opposite the perpendicular.

$$\frac{R \times \text{Sin. } x' = \text{Co. Sin. P.} \times \text{Co. Sin. A.} \therefore x' = \frac{\text{Co. Sin. P.} \times \text{Co. Sin. A.}}{R}.$$

Or by logarithms,  $\text{Log. Sin. } x' = \text{Log. Co. Sin. P.} + \text{Log. Co. Sin. A.} - \text{Log. R.}$   
 $\text{Log. Sin. } x' = 9.7585913 + 9.8546465 - 10. \text{Log. Sin.}$



$x' = 9.6132378 \therefore x' = 24^\circ 13' 55''$ , the complement of which is the required angle =  $65^\circ 46' 5''$ .

2nd. If the perpendicular be  $60^\circ$ , and the

angle opposite the base  $30^\circ$ , require the angle opposite the perpendicular?

3rd. Require the angle opposite the perpendicular, when the angle opposite the base is  $= 45^\circ$ , and the perpendicular  $35^\circ$ ?

Then find  $x''$ , which is the complement of the hypotenuse.

$$R. \times \text{Sin. A.} = \text{Tan. } x'' \times \text{Tan. P.} \quad \text{Tan. } x'' = \frac{R. \times \text{Sin. A.}}{\text{Tan. P.}} \text{ or Log. Tan.}$$


$$x'' = \text{Log. R.} + \text{Log. Sin. A.}$$

$$- \text{Log. Tan. P.} \quad \text{Log. Tan.}$$

$$x'' = 10. + 9.8441979 - 10.$$

$$1547732. \quad \text{Log. Tan. } x'' =$$

$$9.6854247 \therefore x'' = 26^\circ 2' 53''.$$



Then from  $90^\circ \quad 0' \quad 0''$

Take  $26^\circ \quad 2' \quad 53''$

$\hline 63^\circ \quad 57' \quad 7'' = \text{the hypotenuse.}$

2. Given the perpendicular  $= 98^\circ$ , and the angle opposite the base  $= 32^\circ$ , to find the hypotenuse.

## CASE V.

---

WHEN THE TWO LEGS ARE GIVEN  
TO FIND THE HYPOTHENUSE AND  
ANGLES.

To find  $x$ , which represents the complement of the hypotenuse, from whence the hypotenuse is easily known.

$$R. \times \text{Sin. } x = \text{Co. Sin. } P. \times$$

$$\text{Co. Sin. } B. \text{ whence the}$$


$$\text{Sin. } x = \frac{\text{Co. Sin. } P. \times \text{Co. Sin. } B.}{R.}; \text{ or } \text{Log. Sin. } x$$

$$= \text{Log. Co. Sin. } P. + \text{Log. Co. Sin. } B. - \text{Log. } R.$$

$$\text{That is } \text{Log. Sin. } x = 9.7585913 + 9.8842540 -$$

$$10 = 9.6428453 = \text{Log. Co. Sin. } x. \therefore = 26^\circ 3' 53''.$$

$$\text{Consequently the hypotenuse} = 63^\circ 56' 7''.$$




2. Given the base =  $61^\circ$ , and the perpendicular =  $80^\circ$ , require the hypotenuse?

3. Required the hypotenuse when the base is  $90^\circ$ , and the perpendicular  $23^\circ 28'$ ?

4. When the perpendicular is  $37^\circ 42'$ , and the base  $49^\circ 50'$ , what is the hypotenuse?

Then find  $x'$ , which stands in the diagram for the complement of the angle opposite the base.



R.  $\times$  Sin. P. — Tan. B.  
 $\times$  Tan.  $x'$ . Consequently  
 Tan.  $x' = \frac{\text{R.} \times \text{Sin. P.}}{\text{Tan. B.}}$ ; or

Log. Tan.  $x' = \text{Log. R.} + \text{Log. Sin. P.} - \text{Log. Tan. B.}$

Log. Tan.  $x' = 10. + 9.9133645 - 9.9238135 = 9.9895510 = \text{The Log. Tan. of } x' \therefore x' = 44^\circ 18' 39''$  and  $90^\circ - 44^\circ 18' 39'' = 45^\circ 41' 21''$ , the angle required.

2. Given the base, and perpendicular, =  $57^\circ 15'$  and  $39^\circ 30'$  respectively, required the angle opposite the base?


## CASE VI.

---

GIVEN THE TWO ANGLES, TO FIND THE OTHER PARTS OF THE RIGHT-ANGLED SPHERICAL TRIANGLE, OBSERVING THAT THE RIGHT ANGLE IS ALWAYS KNOWN.

Let the angle opposite the base be  $= 45^{\circ} 41' 21''$  then  $A.$  is  $= 44^{\circ} 18' 39''$

its complement; also let the angle opposite the perpendicular  $= 65^{\circ} 46' 5''$  its complement  $= 24^{\circ} 13' 55'' = a$ . The values of  $x$ ,  $x'$ ,  $x''$ , may be found in



the same manner as in the preceding cases.

The multiplicity of Examples being only given to exercise the student.

OF OBLIQUE-ANGLED SPHERICAL  
TRIANGLE TRIGONOMETRY.

---

Every oblique-angled spherical triangle being composed of six parts,—namely, three sides, and three angles; any three being given, the remaining three may be found.

- Let there be given
- I. Two sides and an opposite angle.
  - II. Two sides and an included angle.
  - III. Two angles and an opposite side.
  - IV. Two angles and an included side.
  - V. Three sides.
  - VI. Three angles.


The four first cases may be solved by the formula, page 6, and the other two must have separate rules.

## CASE I.

---

GIVEN TWO SIDES AND AN ANGLE, OPPOSITE ONE OF THEM TO FIND THE REMAINING PARTS OF THE SPHERICAL TRIANGLE.


Given the sides  $AB$ ,  $AC$ , and either of the angles  $ABC$ , or  $ACB$ , to find the side  $BC$  and the remaining angles?



We must suppose a perpendicular  $AD$  to be let fall in such a manner, that two of the known parts of the spherical triangle may be in the right-angled spherical triangle  $BAD$  or  $DAC$ .



Then it is easily observed, that a figure  $HAH'Ba$ , which is double of the one given in right-angled spherical triangle Trigonometry, will answer for oblique-angled spherical triangle Trigonometry.



R. = The Radius.

P. = The perpendicular AD.

B. = The base BD.

$a$ . = The complement of the angle B.

H. = The complement of the hypotenuse AB.

A. = The complement of the angle BAB.

$B'.$  = The base  $CD$ .

$a'.$  = The complement of the angle  $ACD$ .

$H'.$  = The complement of the hypotenuse  $AC$ .



$A'.$  = The complement of the angle  $CAD$ .

In this case the solution is evident, only to keep in mind that when the perpendicular falls outside the base, as in fig., the (1)  $A \hookrightarrow A'$  is the angle  $BAC$ , and  $B \hookrightarrow B'$  is the base  $BC$ , and the Co.  $a'$  is the supplement of the angle  $ACB$ , and not the angle itself; but when the perpendicular falls within the spherical triangle as in fig. (2)  $\text{Co. } A + \text{Co. } A'$  is the angle  $BAC$ ,  $B + B$  is the base  $BC$ , and  $\text{Co. } a'$  is the angle  $ACB$ .

## CASE II.

---

GIVEN ANY TWO SIDES AND THEIR INCLUDED ANGLE OF AN OBLIQUE SPHERICAL TRIANGLE, TO FIND THE REMAINING PARTS OF THE TRIANGLE.




(2)

Given  $H$ ,(or  $AB$ ), $B \leftarrow B'$  or $B + B'$  or $(BC)$ ;

also the

angle



 $ABC$ , tofind the side  $AC$ , and the angles  $BAC$  and  $BCA$ .

It must be observed that the perpendicular arc  $AD$  must be let fall from the point  $A$ , on the base  $BC$ , or  $BC$ , produced; it may be also let fall from the point  $C$ , on the great circle  $AB$ , or  $AB$ , produced.

From what has been said, the solution of the third and fourth cases will easily be arranged and solved; and it may not be amiss to state here again, that  $A \leftarrow A'$  is equal to the angle  $BAC$ , when the perpendicular arc  $AD$  falls outside the base  $BC$ , and  $BC = B \leftarrow B'$ ; but when the arc  $AD$  falls inside of the triangle

Co.  $A + \text{Co. } A'$ , is = the angle  $BAC$  and  $B + B' = BC$  the base.

It happens that the angle  $ACD$  is the supplement of the angle  $ACB$ , when the perpendicular falls outside the triangle, or Co.  $a'$  must be taken from 180 degrees, to give the angle  $ACB$ .



In order that this system may be more simplified, the diagram given in the work may be imagined as placed upon the fingers of the left hand, as in the annexed scheme :


The thumb is marked B, and represents the base.

The little finger is marked P, and represents the perpendicular.


The ring-finger is marked A, and represents the complement of the angle opposite the base.

The middle finger is marked H, and represents the complement of the hypotenuse.

And the finger next the thumb represents the complement of the angle opposite the perpendicular, and is marked  $a$ . Thus the diagram, so familiar in the work, may be conceived to exist on the fingers of our hand, and in short may be slightly marked, until their positions are known.



From what has been said of right-angle tri-angled spherical trigonometry, with



respect to the left hand, we can easily conceive the mode of arranging both hands (as in the annexed plate) for oblique-angled spherical triangle trigonometry.



# DEMONSTRATION

OF THE

## FOREGOING FORMULÆ.

The formulæ given page 6 are demonstrated as follow :—

ABC is any right-angled spherical triangle BC, the base AC, the perpendicular CAB, the angle opposite the base ; AB, the hypothenuse, and ABC, the angle opposite the perpendicular.





It is a well-known property that  $\text{Sin. BC.} : \text{Co. Tan. B.} :: \text{Tan. AC.} : \text{Radius (R.)} \therefore \text{R.} \times \text{Sin. BC.} = \text{Co. Tan. B.} \times \text{Tan. AC.}$

And it is also demonstrable that  $\text{Sin. AB.} : \text{R.} :: \text{Sin. BC.} : \text{Sin. A.} \therefore \text{R.} \times \text{Sin. BC.} = \text{Sin. BA.} \times \text{Sin. A.}$

Secondly, The analogy,  $\text{Co. Sin. B.} : \text{Co. Tan. BA.} :: \text{Tan. BC.} : \text{R.}$  is well known.  $\therefore \text{R.} \times \text{Co. Sin. B.} = \text{Co. Tan. AB.} \times \text{Tan. BC.}$

And  $\text{Co. Sin. AC.} : \text{R.} :: \text{Co. Sin. B.} : \text{Sin. A.} \therefore \text{R.} \times \text{Co. Sin. B.} = \text{Co. Sin. AC.} \times \text{Sin. A.}$

Lastly, By two other well-known analogies,  $\text{Co. Sin. BA.} : \text{Co. Tan. B.} :: \text{Co. Tan. A.} : \text{R.}$  and  $\text{Co. Sin. AC.} : \text{R.} :: \text{Co. Sin. BA.} : \text{Co. Sin. BC.}$

$\therefore \text{R.} \times \text{Co. Sin. AB.} = \text{Co. Tan. B.} \times \text{Co. Tan. A.}$  and  $\text{R.} \times \text{Co. Sin. AB.} = \text{Co. Sin. AC.} \times \text{Co. Sin. BC.}$

Consequently, the general formula page 6 is right.

## CASE V.

---

GIVEN THE THREE SIDES TO FIND THE  
THREE ANGLES.


RULE.

From half the sum of the three sides subtract each of the two sides which contain the required angle.

Then add together the Sines of those two remainders, and the Co-arcs of the Sines of the sides which contain the angle. Half the sum of those four Logs. will give the Sine of half the required angle.

EXAMPLE.

$$\begin{array}{r}
 AB. = 79^{\circ} 17' 14'' \\
 BC. = 100^{\circ} 0' 0'' \\
 AC. = 58^{\circ} 0' 0'' \\
 \hline
 2) 247^{\circ} 17' 14''
 \end{array}
 \left. \vphantom{\begin{array}{r} AB. \\ BC. \\ AC. \\ 2) \end{array}} \right\} \begin{array}{l} \text{To find the} \\ \text{angles.} \end{array}$$



From  $123^{\circ} 38' 37'' =$  the  $\frac{1}{2}$  Sum

Take  $79^{\circ} 17' 14''$

1st rem<sup>r</sup>.  $44^{\circ} 21' 23''$

The Sine of which is  $= 9.8445513$

Again, from  $123^{\circ} 38' 37''$

Take  $58^{\circ} 0' 0''$

2nd rem<sup>r</sup>.  $65^{\circ} 38' 37''$

The Sine of which is  $= 9.9595173$

The Co. Arc. Sin. of  $58^{\circ} 0' 0'' = 0.0715795$

The Co. Arc. Sin. of  $79^{\circ} 17' 14'' = 0.0076359$

2) 19.8832840

9.9416420

Sine  $60^{\circ} 57' 28''$

2

$121^{\circ} 54' 56'' =$  the angle A.

## C A S E V I.

---

WHEN THE THREE ANGLES ARE GIVEN  
TO FIND THE THREE SIDES.

### R U L E.

From half the sum of the three angles subtract each of the angles next the required side.


Then add together the Co. Sines of those two remainders, and the Co-Arc of the Sines of each of the adjoining angles. Half the sum of those four Logs. will give the Co. Sin. of half the required side.

### E X A M P L E.

$$\begin{array}{r}
 \text{Given } A. = 121^{\circ} 54' 56'' \\
 \quad B. = 50^{\circ} 0' 0'' \\
 \quad C. = 62^{\circ} 34' 6'' \\
 \hline
 2) 234^{\circ} 29' 2''
 \end{array}$$

}

To find the  
sides.



Then from  $117^{\circ} 14' 31'' =$  the  $\frac{1}{2}$  Sum

Take  $62^{\circ} 34' 6'' =$  the angle C.

1st remain<sup>r</sup>.  $54^{\circ} 40' 25''$  Co. Sin. = 9.7621032

Again, from  $117^{\circ} 14' 31''$

Take  $50^{\circ} 0' 0''$

2nd remain<sup>r</sup>.  $67^{\circ} 14' 31''$  Co. Sin. = 9.5875321

The Co. Arc Sin.  $50^{\circ} 0' 0'' = 0.1157460$

The Co. Arc Sin.  $62^{\circ} 34' 6'' = 0.0518018$

2) 19.5171831

9.7585915

Co. Sin.  $55^{\circ} 0'$

2

$110^{\circ} 0'$  the side BC required.

---

THE END.

---





QR

535

297

1835



